

RevertAid™ Premium
First Strand cDNA Synthesis Kit
#K1651, #K1652

CERTIFICATE OF ANALYSIS

RevertAid™ Premium First Strand cDNA Synthesis Kit

#__
Lot

QUALITY CONTROL

Functionally tested in RT-PCR using 100fg of control GAPDH RNA and GAPDH control primers generated a prominent 496 bp product on 1% agarose gel after ethidium bromide staining.

Quality authorized by:

 Jurgita Zilinskiene

CONTENTS	page
COMPONENTS OF THE KIT	2
STORAGE	2
DESCRIPTION.....	2
IMPORTANT NOTES.....	3
CONTROL REACTIONS for RT-PCR/RT-qPCR.....	4
PROTOCOLS.....	4
RT-PCR.....	5
RT-qPCR.....	6
cDNA Synthesis for Cloning	7
TROUBLESHOOTING	8
RELATED PRODUCTS.....	9
REFERENCES.....	10

COMPONENTS OF THE KIT

RevertAid™ Premium First Strand cDNA Synthesis Kit	20 rxns #K1651	100 rxns #K1652
RevertAid™ Premium Enzyme Mix	25 µl	120 µl
5X RT Buffer 250 mM Tris-HCl (pH 8.3 at 25°C), 375 mM KCl, 15 mM MgCl ₂ , 50 mM DTT	150 µl	500 µl
10 mM dNTP Mix	50 µl	250 µl
Oligo(dT) ₁₈ Primer 100 µM, 0.5 µg/µl	25 µl	120 µl
Random Hexamer Primer 100 µM, 0.2 µg/µl	25 µl	120 µl
Water, nuclease-free	1.25 ml	2x1.25 ml

STORAGE

All kit components should be stored at -20°C.

DESCRIPTION

The RevertAid™ Premium First Strand cDNA Synthesis Kit is a complete system for efficient synthesis of first strand cDNA. The kit uses the RevertAid™ Premium Reverse Transcriptase (RT), an advanced enzyme derived by *in vitro* evolution of M-MuLV RT. The enzyme features the highest thermostability among the derivatives of M-MuLV RT and lacks the RNase H activity. The RevertAid™ Premium First Strand cDNA Synthesis Kit is capable of synthesizing cDNA up to 20 kb from a wide range of starting RNA amounts (1 pg to 5 µg) at elevated temperatures (50-60°C) superceeding other systems in ability to prepare full length cDNA. Due to the increased synthesis rate reaction can be completed in 30 min.

The RevertAid™ Premium Enzyme Mix contains RevertAid™ Premium Reverse Transcriptase and RiboLock™ RNase Inhibitor. The recombinant RiboLock™ RNase Inhibitor effectively protects RNA from degradation by RNases A, B and C at temperatures up to 55°C.

The kit is supplied with both oligo(dT)₁₈ and random hexamer primers. Random hexamer primers bind non-specifically and are used to synthesize cDNA from all RNAs in total RNA population. The oligo(dT)₁₈ primer selectively anneals to the 3'-end of poly(A) RNA, synthesizing cDNA only from poly(A) tailed mRNA. Gene-specific primers may also be used with the kit to prime synthesis from a specified sequence.

The synthesized cDNA can be used directly in PCR with a variety of thermostable DNA polymerases, in qPCR with Maxima™ qPCR Master Mixes or in second strand cDNA synthesis.

IMPORTANT NOTES

Avoiding ribonuclease contamination

RNA purity and integrity is essential for synthesis of full-length cDNA. RNA can be degraded by RNase A, which is a highly stable contaminant found in any laboratory environment. All components of the kit have been rigorously tested to ensure that they are RNase free. To prevent contamination both the laboratory environment and all prepared solutions must be free of RNases.

General recommendations to avoid RNase contamination:

- Use certified nuclease-free labware or DEPC-treat all tubes and pipette tips to be used in cDNA synthesis.
- Wear gloves when handling RNA and all reagents, as skin is a common source of RNases. Change gloves frequently.
- Use RNase-free reagents, including high quality water (e.g., Water, nuclease-free, #R0581).
- Keep the kit components tightly sealed when not in use. Keep all tubes tightly closed during the reverse transcription reaction.

Template RNA

Total cellular RNA isolated by standard methods is suitable for use with the kit. Purified RNA must be free of salts, metal ions, ethanol and phenol to avoid inhibiting the cDNA synthesis reaction.

For RT-PCR applications, template RNA must be free of DNA contamination. Prior to cDNA synthesis, RNA can be treated with DNase I, RNase-free (#EN0521) to remove trace amounts of DNA. Always perform a RT-minus negative control reaction, which includes all components for RT-PCR except the RevertAid™ Premium Enzyme Mix.

RNA sample quality

Assess RNA integrity prior to cDNA synthesis. Total eukaryotic RNA can be analyzed by agarose gel electrophoresis followed by ethidium bromide staining. The RNA is considered to be intact, if both 18S and 28S rRNA appear as sharp bands after electrophoresis of total RNA. The 28S rRNA band should be approximately twice as intense as the 18S rRNA. Any smearing of rRNA bands is an indication of degraded mRNA. If this occurs, a new sample of total RNA should be prepared. Alternatively, total RNA can be analyzed using a bioanalyzer (e.g., Agilent 2100) which provides quantitative information about the general state of the RNA sample, the RNA integrity number (2). A reference gene/target gene 3':5' integrity assay (3) can also be used to determine the integrity of the RNA sample.

RNA quantity

- Use 1 pg - 5 µg of total RNA, 0.1 pg - 500 ng of poly(A) mRNA or 0.01 pg – 500 ng of specific RNA transcript to generate first strand cDNA as the initial step of a two-step RT-PCR protocol.
- Use 1 µg of isolated mRNA to generate first strand cDNA for second-strand synthesis and subsequent cloning reactions.

Primers

Synthesis of first strand cDNA can be primed with either oligo(dT)₁₈ primer, random primers or gene-specific primers.

Oligo(dT)₁₈ primes cDNA synthesis from the poly(A) tail present at the 3'-end of eukaryotic mRNA. Random primers initiate cDNA synthesis from the total RNA population (rRNA and mRNA). Therefore, using random primers for first strand synthesis results in a greater complexity of the generated cDNA compared with the oligo(dT)₁₈ primer. As a consequence, the sensitivity and specificity of subsequent PCR reactions may be reduced. However, there are several applications where it is beneficial to use random primers, such as cDNA synthesis using mRNAs without a poly(A) tail, or cDNA synthesis using poly(A)-enriched RNA samples. Gene-specific primers are used to synthesize specific cDNA from a pool of total RNA or mRNA and must be obtained by the user.

CONTROL REACTIONS for RT-PCR / RT-qPCR

The following negative control reactions should be used to verify the results of the first strand cDNA synthesis.

- **Reverse transcriptase minus (RT-) negative control** is important in RT-PCR and RT-qPCR reactions to assess for genomic DNA contamination of the RNA sample. The control RT- reaction should contain all reagents for the reverse transcription reaction except for the RevertAid™ Premium Enzyme Mix.
- **No template control (NTC)** is important to assess for reagent contamination. The NTC reaction should contain all reagents necessary for the reverse transcription reaction except for the RNA template.

PROTOCOLS

Please read the IMPORTANT NOTES section of the manual (p.3-4) before starting.

RT-PCR

I. First Strand cDNA Synthesis

After thawing, mix and briefly centrifuge the components of the kit. Store on ice.

1. Add the following reagents into a sterile, RNase free tube on ice in the indicated order:

Template RNA	total RNA or poly(A) mRNA or specific RNA	1 pg - 5 µg 0.1 pg – 500 ng 0.01 pg - 500 ng
Primer	oligo (dT) ₁₈ primer or random hexamer primer or gene-specific primer	1 µl (100 pmol) 1 µl (100 pmol) 2-20 pmol
10 mM dNTP Mix		1 µl (0.5 mM final concentration)
Water, nuclease-free		to 15 µl

2. *Optional.* If the RNA template is GC-rich or is known to contain secondary structures, mix gently, centrifuge briefly and incubate at 65°C for 5 min. Chill on ice, briefly centrifuge again and place on ice.
3. Add the following components in the indicated order:

5X RT Buffer	4 µl
RevertAid™ Premium Enzyme Mix	1 µl
Total volume	20 µl

Mix gently and centrifuge.

4. Incubate:
 - if an oligo(dT)₁₈ primer or gene-specific primer is used, incubate for 30 min at 50°C.
 - if a random hexamer primer is used, incubate for 10 min at 25°C followed by 30 min at 50°C.
 - for transcription of GC-rich RNA, the reaction temperature can be increased to 60°C.
5. Terminate the reaction by heating at 85°C for 5 minutes.

The reaction product of the first strand cDNA synthesis can be used directly in PCR, or stored at -20°C for up to one week. For longer storage, -70°C is recommended. Avoid freeze/thaw cycles of the cDNA.

II. PCR

The product of the first strand cDNA synthesis reaction can be used directly in PCR. The volume of the first strand cDNA synthesis reaction mixture should not comprise more than 1/10 of the total PCR reaction volume. Normally, 2 µl of the first strand cDNA synthesis reaction mixture is used as template for subsequent PCR in a 25 µl total volume. *Taq* DNA polymerase, PCR Master Mix (2X) or Maxima™ Hot Start PCR Master Mix (2X) can be used to amplify fragments less than 3 kb. DreamTaq™ DNA polymerase is suitable for amplification of longer fragments up to 6 kb. The Long PCR Enzyme Mix and High Fidelity PCR Enzyme Mix are recommended to generate amplicons up to 20 kb. See related products on p.10 for ordering information.

RT-qPCR

I. First Strand cDNA Synthesis

The following protocol is optimized to generate first-strand cDNA for use in two-step RT-qPCR. After thawing, mix and briefly centrifuge the components of the kit. Store on ice.

1. Add the following reagents into a sterile, RNase free tube on ice in the indicated order:

Template RNA	total RNA or poly(A) mRNA or specific RNA	1 pg - 5 µg 0.1 pg – 500 ng 0.01 pg - 500 ng
Primers	oligo (dT) ₁₈ primer and random hexamer primer or gene-specific primer	0.25 µl (25 pmol) 0.25 µl (25 pmol) 2-20 pmol
10 mM dNTP Mix		1 µl
Water, nuclease-free		to 15 µl

2. *Optional.* If the RNA template is GC-rich or is known to contain secondary structures, mix gently, centrifuge briefly and incubate at 65°C for 5 min. Chill on ice, briefly centrifuge again and place on ice.
6. Add the following components to the reaction tube in the indicated order:

5X RT Buffer	4 µl
RevertAid™ Premium Enzyme Mix	1 µl
Total volume	20 µl

Mix gently and centrifuge.

3. Incubate for 10 min at 25°C followed by 15 min at 50°C.
4. Terminate the reaction by heating at 85°C for 5 minutes.

Note. For RNA template quantities greater than 1 µg, prolong the reaction time to 30 min. For RNA templates that are GC-rich or have a large amount of secondary structure, the reaction temperature can be increased to 60°C.

The reverse transcription reaction product can be used directly in qPCR, or stored at -20°C for up to one week. For longer storage, -70°C is recommended. Avoid freeze/thaw cycles of the cDNA.

II. qPCR

The product of the first strand cDNA synthesis reaction can be used directly in qPCR. The volume of first strand cDNA synthesis reaction mixture should not comprise more than 1/10 of the total PCR reaction volume. Normally, 2 µl of the RT mixture is used as template for subsequent qPCR in 25 µl total volume. RevertAid Premium™ First Strand cDNA Synthesis Kit is optimized for use with Maxima™ qPCR master mixes (see Related Products).

cDNA Synthesis for Cloning

I. First strand cDNA synthesis

Mix and briefly centrifuge all components after thawing, keep on ice.

1. Add the following reagents into a sterile, nuclease-free tube on ice in the indicated order:

Template RNA	poly(A) mRNA or specific RNA	1 µg 0.5-1 µg
Primer	oligo (dT) ₁₈ primer or random hexamer primer or gene-specific primer	1 µl 1 µl 100 pmol
10 mM dNTP Mix		1 µl
Water, nuclease-free		to 15 µl

2. *Optional.* If the RNA template is GC-rich or contains secondary structures, mix gently, centrifuge briefly and incubate at 65°C for 5 min. Chill on ice, spin down and place the vial back on ice.
3. Add the following components to the reaction tube in the indicated order:

5X RT Buffer	4 µl
RevertAid™ Premium Enzyme Mix	1 µl
Total volume	20 µl

Mix gently and centrifuge.

4. Incubate:
 - if an oligo(dT)₁₈ primer or gene-specific primer is used, incubate for 30 min at 50°C.
 - if a random hexamer primer is used, incubate for 10 min at 25°C followed by 30 min at 50°C.
 - for transcription of GC-rich RNA, the reaction temperature can be increased to 60°C.
5. Terminate the reaction by heating at 85°C for 5 min.

The reverse transcription reaction product can be used immediately in second strand cDNA synthesis reactions or stored at -20°C for up to one week. For longer storage, -70°C is recommended.

II. Second strand cDNA synthesis

1. Add the following reagents to 20 µl of the first strand cDNA synthesis reaction mixture on ice:

10X reaction buffer for DNA Polymerase I	8 µl
RNase H (#EN0201)	0.2 µl (1 u)
DNA Polymerase I (#EP0041)	3 µl (30 u)
Water, nuclease-free (#R0581)	to 100 µl
Total volume	100 µl

2. Gently vortex and briefly centrifuge.
3. Incubate at 15°C for 2 hours. Do not let the temperature to rise above 15°C.
4. Add 2.5 µl (12.5 u) of T4 DNA Polymerase (#EP0061) and incubate at 15°C for 5 min.
5. Terminate the reaction by adding 5 µl of 0.5 M EDTA, pH 8.0 (#R1021).

Purify blunt-end cDNA using the GeneJET™ PCR Purification Kit (#K0701) or by phenol/chloroform extraction and use for further cloning-related procedures e.g., adapter ligation, phosphorylation, size fractionation, ligation and transformation.

TROUBLESHOOTING

Problem	Cause and Solution
<p>Low yield or no product in RT-PCR or RT-qPCR</p>	<p>Poor integrity of RNA template. RNA purity and integrity is essential for synthesis and quantification of cDNA. Always assess the integrity of RNA prior to cDNA synthesis (see page 3). Use freshly prepared RNA. Multiple freeze/thaw cycles of the RNA sample and synthesized cDNA is not recommended. Follow general recommendations to avoid RNase contamination (page 3) and discard low quality RNA.</p> <p>Low template purity (inhibitors in RNA sample). Trace amounts of agents used in RNA purification protocols may remain in solution and inhibit first strand synthesis, e.g., SDS, EDTA, guanidine salts, phosphate, pyrophosphate, polyamines, spermidine. To remove trace contaminants, re-precipitate the RNA with ethanol and wash the pellet with 75% ethanol.</p> <p>Insufficient template quantity. Increase the amount of RNA template in the first strand reaction to the recommended level. Following DNase I treatment, terminate the reaction by heat inactivation of the enzyme in the presence of EDTA (to bind Mg²⁺).</p> <p>GC rich template. If the RNA template is GC rich or is known to contain secondary structures, the temperature of the reverse transcription reaction can be increased up to 60°C.</p> <p>Excess amount of cDNA in qPCR/PCR. Decrease amount of cDNA synthesis reaction in qPCR/PCR. The volume of the cDNA synthesis reaction mixture should not exceed 10% of the final PCR reaction mixture.</p>
<p>No product in long RT-PCR (>5 kb)</p>	<p>Suboptimal priming. Use oligo (dT)₁₈ primer or gene specific primer. If random primers are used, reduce the amount of random primers to 25 pmol per 20 µl reaction mixture.</p>
<p>RT-PCR product longer than expected</p>	<p>RNA template is contaminated with DNA. Amplification of genomic DNA containing introns. To avoid amplification of genomic DNA, design PCR primers on exon-exon boundaries or perform DNase I digestion prior reverse transcription.</p>
<p>RT-PCR product in RT-minus control</p>	<p>RNA template is contaminated with DNA. The presence of a PCR product in the negative control (RT-) reaction indicates that the reaction is contaminated with DNA. To avoid amplification of genomic DNA, design PCR primers on exon-exon boundaries or perform DNase I digestion prior reverse transcription.</p>

RELATED PRODUCTS

Product	Amount	Catalog #
Maxima™ SYBR Green/ROX qPCR Master Mix (2X)	2 x 1.25 ml	K0221
	10 x 1.25 ml	K0222
	4 x 12.5 ml	K0223
Maxima™ SYBR Green/Fluorescein qPCR Master Mix (2X)	2 x 1.25 ml	K0241
	10 x 1.25 ml	K0242
Maxima™ SYBR Green qPCR Master Mix (2X), ROX Solution provided	2 x 1.25 ml	K0251
	10 x 1.25 ml	K0252
Maxima™ Probe/ROX qPCR Master Mix (2X)	2 x 1.25 ml	K0231
	10 x 1.25 ml	K0232
	4 x 12.5 ml	K0233
Maxima™ Probe qPCR Master Mix (2X), ROX Solution provided	2 x 1.25 ml	K0261
	10 x 1.25 ml	K0262
Maxima™ Hot Start <i>Taq</i> DNA Polymerase	100 u	EP0601
	500 u	EP0602
	5 x 500 u	EP0603
TrueStart™ <i>Taq</i> DNA Polymerase	100 u	EP0611
	500 u	EP0612
	5 x 500 u	EP0613
Maxima™ Hot Start PCR Master Mix (2X)	2 x 1.25 ml	K1051
	10 x 1.25 ml	K1052
DreamTaq™ DNA Polymerase	200 u	EP0701
	500 u	EP0702
	5 x 500 u	EP0703
<i>Taq</i> DNA Polymerase, recombinant	100 u	EP0401
	500 u	EP0402
PCR Master Mix (2X)	4 x 1.25 ml	K0171
	20 x 1.25 ml	K0172
<i>Pfu</i> DNA Polymerase, recombinant*	100 u	EP0501
	500 u	EP0502
Long PCR Enzyme Mix*	100 u	K0181
	500 u	K0182
High Fidelity PCR Enzyme Mix*	100 u	K0191
	500 u	K0192
dNTP Mix, 10 mM each	0.2 ml	R0191
	1 ml	R0192
	5 x 1 ml	R0193
Water, nuclease-free	4 x 1.25 ml	R0581
	30 ml	R0582
0.5 M EDTA, pH 8.0	5 x 1 ml	R1021
DNA Polymerase I	500 u	EP0041
	5 x 500 u	EP0042
RNase H	100 u	EN0201
	500 u	EN0202
DNase I, RNase-free	1000 u	EN0521
GeneJET™ PCR Cloning Kit	50 preps	K0701
	250 preps	K0702
CloneJET™ PCR Purification Kit	20 rxns	K1231
	40 rxns	K1232

*Not available in the USA.

References

1. Wiame, I., et al., Irreversible heat inactivation of DNaseI without RNA degradation, *BioTechniques*, 29, 252-256, 2000.
2. Fleige, S., Pfaffl, M.W., RNA integrity and the effect on the real-time qRT-PCR performance, *Mol. Aspects Med.*, 27, 126-139, 2006.
3. Nolan, T., et al., Quantification of mRNA using real-time RT-PCR, *Nat. Protoc.*, 1, 1559-1582, 2006.

This product or the use of this product is covered by international patent application WO/2009/125006. The purchase of this product includes a non-transferable license to use this product for the purchaser's internal research. All other commercial uses of this product, including without limitation product use for diagnostic purposes, resale of product in the original or any modified form or product use in providing commercial services require a separate license from Fermentas. For further information on obtaining licenses please contact Fermentas at info@fermentas.com.

Maxima, RevertAid, RiboLock, DreamTaq, TrueStart, GeneJET and CloneJET are trademarks of Fermentas.

PRODUCT USE LIMITATION.

This product is developed, designed and sold exclusively for research purposes and *in vitro* use only. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.fermentas.com for Material Safety Data Sheet of the product.

